

Compositeness of exotic hadrons with decay and coupled-channel effects



arXiv:2303.07038 [hep-ph]



Tomona Kinugawa

Tetsuo Hyodo

Department of Physics, Tokyo Metropolitan University
May 26th, seminar at Tokyo Tech.

Contents

1. Introduction

compositeness

2. Compositeness of shallow (quasi) bound states

- single-channel scattering model

T. Kinugawa and T. Hyodo, arXiv:2303.07038 [hep-ph].

with decay channel effect

with coupled channel effect

- application to T_{cc} & $X(3872)$

3. New interpretation of compositeness of resonances

- definition of interpretation

T. Kinugawa and T. Hyodo, in preparation

- application to resonances with narrow decay width

4. Summary

Contents

1. Introduction

compositeness

2. Compositeness of shallow (quasi) bound states

T. Kinugawa and T. Hyodo, arXiv:2303.07038 [hep-ph].

- single-channel scattering model

with decay channel effect

with coupled channel effect

- application to T_{cc} & $X(3872)$

3. New interpretation of compositeness of resonances

T. Kinugawa and T. Hyodo, in preparation


- definition of interpretation

- application to resonances with narrow decay width

4. Summary

Near-threshold exotic hadrons


$$X(3872) \rightarrow \pi^+ \pi^- J/\psi$$




S. K. Choi *et al.* (Belle), Phys. Rev. Lett. **91**, 262001 (2003).

internal structure?



exotic hadron
 $\neq qqq$ or $q\bar{q}$



multiquarks
hadronic molecules



LHCb Collaboration, Nature Phys. **18** (2022) no.7, 751-754;
LHCb Collaboration, Nat. Commun. **13** 3351 (2022).





Compositeness

○ definition

hadron wavefunction

$$|\Psi\rangle = \sqrt{X} |\text{hadronic molecule}\rangle + \sqrt{1-X} |\text{others}\rangle$$

compositeness



elementarity

T. Hyodo, Int. J. Mod. Phys. A 28, 1330045 (2013).

* $0 \leq X \leq 1 \longrightarrow X > 0.5 \Leftrightarrow \text{composite dominant}$

$X < 0.5 \Leftrightarrow \text{elementary dominant}$

○ advantage

- **quantitative** analysis of internal structure of bound states
- applicable not only to hadrons but also to nuclei and atoms

History of compositeness

- Weinberg's work (1960s) Weinberg, S. Phys. Rev. 137, 672–678 (1965) etc.
deuteron is not an elementary particle ← weak-binding relation
- application to exotic hadrons (2000s-)
 - “compositeness”
 - generalization to unstable states
 - with spectral function V. Baru, J. Haidenbauer, C. Hanhart, Y. Kalashnikova, A.E. Kudryavtsev, Phys. Lett. B 586, 53–61 (2004) etc.
 - with effective range expansion T. Hyodo, Phys. Rev. Lett. **111**, 132002 (2013) etc.
 - with effective field theory Y. Kamiya and T. Hyodo, PTEP 2017, 023D02 (2017) etc.
 - application to ...
 - $f_0(980)$, $a_0(980)$ Y. Kamiya and T. Hyodo, PTEP 2017, Phys. Rev. C 93, 035203 (2016);
T. Sekihara, S. Kumano, Phys. Rev. D 92, 034010 (2015) etc.
 - $\Lambda(1405)$ T. Sekihara, T. Hyodo, Phys. Rev. C 87, 045202 (2013) ;
Z.H. Guo, J.A. Oller, Phys. Rev. D 93, 096001 (2016) etc.
 - nuclei & atomic systems T. Kinugawa, T. Hyodo, Phys. Rev. C 106, 015205 (2022) etc.

Compositeness

● model calculation

T. Hyodo, D. Jido, and A. Hosaka, Phys. Rev. C **85**, 015201 (2012);
 F. Aceti and E. Oset, Phys. Rev. D **86**, 014012 (2012).

$$T = \frac{1}{V^{-1} - G}$$

V : effective interaction
 G : loop function

residue of scattering amplitude g

$$X = -g^2 G'(E) \Big|_{E=-B} \quad \alpha'(E) = d\alpha/dE$$

$$= \frac{G'(E)}{G'(E) - [V^{-1}(E)]'} \Big|_{E=-B}$$

Y. Kamiya and T. Hyodo, PTEP 2017, 023D02 (2017).

g^2 : model independent $\leftarrow T_{\text{on}}(-B)$ (observable)

$G(E)$: model dependent \leftarrow cutoff dependent

Weak-binding relation

S. Weinberg, Phys. Rev. 137, 672–678 (1965).

$$X = \frac{a_0}{2R - a_0} + \mathcal{O}\left(\frac{R_{\text{typ}}}{R}\right)$$

a_0 : scattering length
 R_{typ} : typical length scale in system
 $R = 1/\sqrt{2\mu B}$

- for weakly bound states, $R \gg R_{\text{typ}}$

compositeness ←— observables (a_0, B)

Y. Li, F.-K. Guo, J.-Y. Pang, and J.-J. Wu, Phys. Rev. D 105, L071502 (2022);
 J. Song, L. R. Dai, and E. Oset, Eur. Phys. J. A 58, 133 (2022);
 M. Albaladejo, J. Nieves, Eur. Phys. J. C 82, 724 (2022);
 T. Kinugawa, T. Hyodo, Phys. Rev. C 106, 015205 (2022).


○ range correction

compositeness of deuteron $X \sim 1.7 > 1$

→ important to consider effective range

- our work : range correction ←— uncertainty estimation

compositeness of deuteron : $0.74 \leq X \leq 1$



Low-energy universality

scattering length $a_0 (\rightarrow \infty)$

\gg typical length scale of system R_{typ}

low-energy universality

E. Braaten and H.-W. Hammer, Phys. Rept. **428**, 259 (2006) ;
F. P. Naidon and S. Endo, Rept. Prog. Phys. **80**, 056001 (2017).

→ length scales are written only by $|a_0|$

- for bound states ?

$$R = 1/\sqrt{2\mu B} : a_0 = R \rightarrow \infty \longrightarrow B \rightarrow 0$$

→ universality holds for **weakly-bound states!**

- compositeness $X = 1$ in $B \rightarrow 0$ limit T. Hyodo, Phys. Rev. C **90**, 055208 (2014) .

→ near threshold states ($B \sim 0$) = composite dominant ?

e.g. ${}^8\text{Be}$, ${}^{12}\text{C}$ Hoyle state → α cluster?

Contents

1. Introduction

compositeness

2. Compositeness of shallow (quasi) bound states

T. Kinugawa and T. Hyodo, arXiv:2303.07038 [hep-ph].

- single-channel scattering model

with decay channel effect

with coupled channel effect

- application to T_{cc} & $X(3872)$

3. New interpretation of compositeness of resonances

T. Kinugawa and T. Hyodo, in preparation


- definition of interpretation

- application to resonances with narrow decay width

4. Summary

Decay & coupled ch. effects

actual exotic hadrons \longrightarrow decay and coupled channel



other ch. than threshold ch. make deviation from $X = 1$

Y. Kamiya and T. Hyodo, PTEP 2017, 023D02 (2017).


In this work, we study those deviations quantitatively!

universality

T_{cc} and $X(3872)$ are shallow-bound states


→ low-energy universality is important!

1. naive expectation : near-threshold states
are composite dominant



2. However, elementary dominant states is realized
with fine tuning

T. Hyodo, Phys. Rev. C **90**, 055208 (2014) ;
C. Hanhart, J. R. Pelaez, and G. Rios, Phys. Lett. B **739**, 375 (2014).



How finely tuning parameter?

In this work, we study fine tuning quantitatively!

Outline of this work

13

T. Kinugawa and T. Hyodo, arXiv:2303.07038 [hep-ph].

Quantitative discussion of universality with simple model
single-channel scattering model (Sec. II)



Introducing decay & coupled-channel effects
decay model and coupled-channel model (Sec. III)



Application to T_{cc} and $X(3872)$
both of decay and coupled-channel model (Sec. IV)

Model

E. Braaten, M. Kusunoki, and D. Zhang, Annals Phys. 323, 1770 (2008).

● single-channel resonance model

$$\mathcal{H}_{\text{free}} = \frac{1}{2m_1} \nabla \psi_1^\dagger \cdot \nabla \psi_1 + \frac{1}{2m_2} \nabla \psi_2^\dagger \cdot \nabla \psi_2 + \frac{1}{2m_\phi} \nabla \phi^\dagger \cdot \nabla \phi + \nu_0 \phi^\dagger \phi,$$



1.

$$\mathcal{H}_{\text{int}} = g_0 (\phi^\dagger \psi_1 \psi_2 + \psi_1^\dagger \psi_2^\dagger \phi).$$

2.

1. single-channel scattering

2. coupling with compact state ϕ



● scattering amplitude

$$V = \frac{g_0^2}{E - \nu_0}, \quad G = -\frac{\mu}{\pi^2} \left[\Lambda + ik \arctan \left(\frac{\Lambda}{-ik} \right) \right]. \quad \Lambda : \text{cutoff}$$

$$T = \frac{1}{V^{-1} - G} \xrightarrow{\longrightarrow} f(k) = -\frac{\mu}{2\pi} \left[\frac{\frac{k^2}{2\mu} - \nu_0}{g_0^2} + \frac{\mu}{\pi^2} \left[\Lambda + ik \arctan \left(\frac{\Lambda}{-ik} \right) \right] \right]^{-1}.$$

Model scales and parameters

- typical energy scale : $E_{\text{typ}} = \Lambda^2/(2\mu)$

- three model parameters g_0, ν_0, Λ

1. calculation with given B

coupling const. g_0 :
$$g_0^2(B, \nu_0, \Lambda) = \frac{\pi^2}{\mu} (B + \nu_0) \left[\Lambda - \kappa \arctan(\Lambda/\kappa) \right]^{-1}$$

\because bound state condition $f^{-1} = 0$ $\kappa = \sqrt{2\mu B}$.

2. use dimensionless quantities with Λ

→ results do not depend on cutoff Λ

3. energy of bare quark state ν_0

varied in the region : $-B/E_{\text{typ}} \leq \nu_0/E_{\text{typ}} \leq 1$

\because to have $g_0^2 \geq 0$ & applicable limit of EFT

Calculation

● compositeness X


scattering amplitude : $T = \frac{1}{V^{-1} - G}$

Y. Kamiya and T. Hyodo,
PTEP 2017, 023D02 (2017).

$$\begin{aligned} \longrightarrow X &= \frac{G'(-B)}{G'(-B) - [V^{-1}(-B)]'}, \quad \alpha'(E) = d\alpha/dE \\ &= \left[1 + \frac{\pi^2 \kappa}{g_0^2 \mu^2} \left(\arctan(\Lambda/\kappa) - \frac{\Lambda/\kappa}{1 + (\Lambda/\kappa)^2} \right)^{-1} \right]^{-1}. \end{aligned}$$

- ν_0 region : $-B/E_{\text{typ}} \leq \nu_0/E_{\text{typ}} \leq 1$


compositeness X as a function of ν_0



\longrightarrow internal structure of bound state?

● X as a function of ν_0/E_{typ} of bound state $B = E_{\text{typ}}$

17




- typical energy scale : $B = E_{\text{typ}} = \Lambda^2/(2\mu)$

- $X > 0.5$ only for 25 % of ν_0 = elementary dominant



∴ bare state origin

● X as a function of ν_0/E_{typ} of bound state $B = 0.01E_{\text{typ}}$ 18



- weakly-bound state : $B = 0.01E_{\text{typ}}$
- $X > 0.5$ for 88 % of ν_0 = composite dominant

∴ low-energy universality !

Effect of decay

● introducing decay effect

- formally : introducing decay channel in lower energy region than binding energy

→ eigenenergy becomes complex

- effectively : coupling const. $g_0 \in \mathbb{C}$! ← this work

$$\mathcal{H}_{\text{int}} = \underline{g_0} (\phi^\dagger \psi_1 \phi_2 + \phi_1^\dagger \psi_2^\dagger \phi).$$

$$E = -B \rightarrow E = -B - \underline{i\Gamma/2}$$


compositeness

$$X \in \mathbb{R} \rightarrow X \in \mathbb{C}$$


$$\tilde{X} = \frac{|X|}{|X| + |1-X|}$$

Effect of decay

$$E = -0.01E_{\text{typ}} - i0.1E_{\text{typ}}$$



$$E = -0.01E_{\text{typ}} - iE_{\text{typ}}$$




- \tilde{X} is suppressed by decay effect

\because threshold ch. component (\tilde{X}) decreases with inclusion of decay ch. component ($1 - \tilde{X}$)


Effect of decay

$$E = -0.01E_{\text{typ}} - i0.1E_{\text{typ}}$$



$$(a) E = -0.01E_{\text{typ}} - i0.1E_{\text{typ}}$$

$$E = -E_{\text{typ}} - i0.1E_{\text{typ}}$$



$$\Gamma = 0.1E_{\text{typ}}$$

$$(c) E = -E_{\text{typ}} - i0.1E_{\text{typ}}$$

compositeness is more suppressed when B is small

- suppression of \tilde{X} is determined by ratio of B to Γ

Effect of coupled channel


- introducing coupled channel $\Psi_1 \Psi_2$

$$|\Psi\rangle = \sqrt{X_1} |\text{threshold ch}\rangle + \sqrt{X_2} |\text{coupled ch}\rangle + \sqrt{1 - (X_1 + X_2)} |\text{others}\rangle$$

$$\mathcal{H}_{\text{free}} = (\text{kinetic terms of } \psi_{1,2}, \Psi_{1,2}, \phi) + \underline{\omega_1 \Psi_1 \Psi_1^\dagger + \omega_2 \Psi_2 \Psi_2^\dagger + \nu_0 \phi^\dagger \phi},$$

$$\mathcal{H}_{\text{int}} = \underline{g_0} (\phi^\dagger \psi_1 \psi_2 + \psi_1^\dagger \psi_2^\dagger \phi + \phi^\dagger \Psi_1 \Psi_2 + \Psi_1^\dagger \Psi_2^\dagger \phi).$$


- threshold energy difference $\Delta\omega = \underline{\omega_1 + \omega_2}$
- ch. 1 couples to ch. 2 through ϕ with same coupling const.



- low-energy universality with coupled-channel effect

$X_1 \sim 1$ (threshold channel)


$X_2 \sim 0$ and $Z \sim 0$ (other channel)




coupled ch. 22

Effect of coupled channel

$$(B, \Delta\omega) = (0.01E_{\text{typ}}, 0.01E_{\text{typ}})$$



$$(B, \Delta\omega) = (0.01E_{\text{typ}}, E_{\text{typ}})$$




- X_1 is suppressed by channel coupling

- \therefore threshold ch. component (X_1) decreases with inclusion of coupled ch. component (X_2)

- $Z = 1 - (X_1 + X_2)$ is stable

Application to T_{cc} and $X(3872)$

● exotic hadron ← decay and coupled channel



LHCb Collaboration, Nat. Commun **13** 3351 (2022).


PDG

● compositeness T. Sekihara, *et. al.*, PRC 93, 035204 (2016).

$$\tilde{X}_j = \frac{|X_j|}{\sum_j |X_j| + |Z|}, \quad (j = 1, 2)$$

\tilde{X}_1 : threshold ch. compositeness
 \tilde{X}_2 : coupled ch. compositeness

Application to T_{cc} and $X(3872)$



- \tilde{X}_2 is not negligible


\therefore coupled ch. contribution (small $\Delta\omega$)

- difference of $\tilde{X}_1 + \tilde{X}_2(\Gamma = 0)$ and $\tilde{X}_1 + \tilde{X}_2$ is too small

→ We can neglect decay contribution

$\therefore \Gamma \ll B$

Application to T_{cc} and $X(3872)$




- difference of $\tilde{X}_1 + \tilde{X}_2(\Gamma = 0)$ and $\tilde{X}_1 + \tilde{X}_2$ is large
 \because large decay width contribution
- \tilde{X}_2 is much smaller than \tilde{X}_1
 \rightarrow coupled ch. effect is small

Application to T_{cc} and $X(3872)$

27


- $\Lambda = 140$ MeV (π meson)



- T_{cc} : $\tilde{X}_1 > 0.5$ for 45 % of ν_0 region
- $X(3872)$: $\tilde{X}_1 > 0.5$ for 59 % of ν_0 region
- coupled ch. effect is more important for T_{cc} than $X(3872)$
- decay effect is more important for $X(3872)$ than T_{cc}

Application to T_{cc} and $X(3872)$

- $\Lambda = 770$ MeV (ρ meson)



- $T_{cc} : \tilde{X}_1 > 0.5$ for 85 % of ν_0 region
- $X(3872) : \tilde{X}_1 > 0.5$ for 87 % of ν_0 region
- typical energy scale E_{typ} is larger
 → states becomes close to universality limit $X \rightarrow 1$
 decay effect : suppressed coupled ch. effect : enhanced

Summary of this part

T. Kinugawa and T. Hyodo,
arXiv:2303.07038 [hep-ph]

- internal structure of exotic hadrons ← EFT & compositeness
- shallow bound state
 - composite dominant even from bare state
fine tuning is necessary to realize elementary dominant state
- decay and coupled channel effects are introduced
 - both decay and coupled ch. effect suppress compositeness
- T_{cc} and $X(3872)$ with decay and coupled ch. effects

T_{cc} : important coupled ch. effect with negligible decay effect

$X(3872)$: important decay effect with negligible coupled ch. effect